MEASURE	AGE	MEASURE COMPONENTS			
ONCOLOGY					
Breast Cancer Screenings (BCS)	Women, ages 50-74	Mammogram: October 1, two years prior to the mea- surement year, through December 31 of the measure- ment year: • CPT/CPT II: 77061-77063, 77065-77067			
Colorectal Cancer Screening(COL)*	Adults, ages 45- 75	 CPT/CPT II: 77061-77063, 77065-77067 Colonoscopy: CPT/CPT II: 44388-44394, 44397, 44401-44408, 45355, 45378-45393,45398 HCPCS: G0105, G0121 Computed tomography (CT) colonography: CPT/CPT II: 74261-74263 FIT-DNA test: CPT/CPT II: 81528. This code is specific to the Cologuard® FIT-DNA test. HCPCS: G0464. This code was retired and replaced with CPT/CPT II code 81528 on January 1, 2016. LOINC: 77353-1, 77354-9 Flexible sigmoidoscopy: CPT/CPT I: 45330-45335, 45337-45342, 45346, 45347, 45349, 45350 HCPCS: G0104 FOBT: CPT/CPT II: 82270, 82274 HCPCS: G0328 LOINC: 12503-9, 12504-7, 14563-1, 14564-9, 14565-6, 2335-8, 27396-1, 27401-9, 27925-7, 27926-5, 29771-3, 56490-6, 56491-4, 57905-2, 58453-2, 80372-6 			

* Allows for telephone visits, e-visits, or virtual check-ins.

HEDIS[®] is a registered trademark of the National Committee for Quality Assurance (NCQA). For more information regarding HEDIS[®] MY 2023, please contact Quality Improvement Dept. at 757-252-8400 or toll-free at 1-844-620-1015.

QCQG.10/21

¹

MEASURE AGE MEASURE COMPONENTS CARE FOR OLDER ADULTS Care for Older Adults Adults, ages 66 Medication Review: CPT/CPT II: 1160F 90863, 99483, (COA) and older 99605, 99606 Medication List: CPT II: 1159F Transition Care Management: CPT 99495, 99496 Functional Status Assessment: CPT: 1170F, 99483 Pain Assessment: CPT/CPT II 1125F, 1126F CARDIOLOGY Controlling High Blood Adults, ages 18-Adequate control is defined as: <140/90 Pressure (CBP)* 85 ICD-10 Codes to identify HTN: 110 **CPT II Codes for BP values:** Systolic Less than 140: 3074F, 3075F Systolic Greater than/Equal to 140: 3077F • Diastolic Less than 80: 3078F Diastolic 80-89: 3079F Diastolic Greater than/Equal to 90: 3080F ICD-10 codes for myalgia/myositis/myopathy: Statin Therapy for Patients Men, ages 21-75 with Cardiovascular Disease • Women, ages 40-G72.0, G72.2, G72.9, M60.80-M60.812, M60.819, (SPC) 75 M60.821, M60.822, M60.829, M60.831, M60.832, M60.839, M60.841, M60.842, M60.849, M60.851, M60.852, M60.859, M60.861, M60.862, M60.869, M60.871, M60.872, M60.879, M60.88, M60.89, M60.9, M62.82, M79.11, M79.10-M79.12, M79.18

* Allows for telephone visits, e-visits, or virtual check-ins.

MEASURE AGE **MEASURE COMPONENTS MUSCULOSKELETAL** Osteoporosis Management Women, ages Bone mineral density tests: in Women Who Had a 67-85 CPT/CPT II: 76977, 77078, 77080, 77081, 77085, Fracture (OMW) 77086 • ICD-10 procedure: BP48ZZ1, BP49ZZ1, BP4GZZ1, BP4HZZ1, BP4LZZ1, BP4MZZ1, BP4NZZ1, BP-4PZZ1, BQ00ZZ1, BQ01ZZ1, BQ03ZZ1, BQ04ZZ1, BR00ZZ1, BR07ZZ1, BR09ZZ1, BR0GZZ1 Osteoporosis medications: HCPCS: J0897, J1740, J3110. J3489 MEDICATION Medication Adherence: Adults, ages 18 At least 2 fills of a designated medication(s) as defined in each measure. The measurement is calculated using **Diabetes Medications** and older (MAD) Proportion of Days Covered (PDC) using Pharmacy Data Encounter (PDE) data. PDC is the number of days the Hypertension (RAS Antagonists) (MAH) member has medication on-hand/is covered by the medi-Cholesterol(Statins) cation based on the dates of prescription fills (numerator) compared to the number of days in the time period (de-(MAC) nominator). DIABETES Adults, ages CPT/CPT II: 67028, 67030, 67031, 67036, Eye Exam for Patients 18-75 67039-67043,67101, 67105, 67107, 67108, with Diabetes (EED) 67110, 67113, 67121, 67141, 67145, 67208, 67210, 67218, 67220, 67221, 67227, 67228, 92002, 92004, 92012, 92014, 92018, 92019, 92134, 92201, 92202, 92225-92228, 92230, 92235, 92240, 92250, 92260, 99203-92205, 99213-99215, 99242-99245, 2022F-2026F, 2033F, 3072F HCPCS: S0620, S0621, S3000

* Allows for telephone visits, e-visits, or virtual check-ins.

MEASURE	AGE	MEASURE COMPONENTS		
DIABETES				
Diabetes	Adults, ages 18-75 with diagnosis of diabetes (type 1 or type 2) or who have been prescribed insulin or hypogly- cemic/anti-hyper- glycemics.	•		
Kidney Health Evaluation for Patients with Diabetes (KED)	Adults, ages 18-85 with diabetes (type 1 and type 2) who received a kidney health evaluation, defined by an estimated glomer- ular filtration rate (eGFR) and a urine albumin-creatinine ratio (uACR)	 eGFR: 80047-80050, 80053, 80069, 82565 uACR: 82043, 82570 		

* Allows for telephone visits, e-visits, or virtual check-ins.

MEASURE	AGE	MEASURE COMPONENTS		
DIABETES				
Hemoglobin A1c Control for Patients with Diabetes (HBD)	Adults, ages 18- 75 with diagnosis of diabetes (type 1 or type 2) or who have been prescribed insulin or hypoglycemic/ anti-hyperglyce- mics.	HbA1c test CPT/CPT II: 83036, 83037 (A1c Testing), 3044F, 3046F, 3051F, 3052F HbA1c level < 8.0: CPT/CPT II 3051F HbA1c level > 9.0: CPT/CPT II 3046F		
Statin Use for Patients with Diabetes (SPD)	Adults, ages 40- 75	ICD-10 codes for myalgia, myositis, myopathy: G72.0, G72.2, G72.9, M60.80-M60.812, M60.819, M60.821, M60.822, M60.829, M60.831, M60.832, M60.839, M60.841, M60.842, M60.849, M60.851, M60.852, M60.859, M60.861, M60.862, M60.869, M60.871, M60.872, M60.879, M60.88, M60.89, M60.9, M62.82, M79.11, M79.12, M79.18		

* Allows for telephone visits, e-visits, or virtual check-ins.